arXiv:1702.00086v1 [math.GT] 31 Jan 2017

RIBBON n-KNOTS WITH ISOMORPHIC QUANDLES

BLAKE K. WINTER

Abstract. Let K, K' be ribbon knottings of $S^{n-1} \times S^{n+2}$, $n \geq 2$. We show that if the knot quandles of these knots are isomorphic, then the ribbon knottings are stably equivalent, in the sense of Nakanishi and Nakagawa, after taking a finite number of connected sums with trivially embedded copies of $S^{n-1} \times S^{n+2}$.

1. Introduction

2. Ribbon Links

Because we will be making reference to ribbon links and their ribbon presentations in further sections, we give here a brief review of these concepts. See [2] for further discussion of ribbon links in arbitrary dimensions.

Remark 1. *Throughout, we will use the notation D^n to represent a closed n-ball.*

Let M be an $(n+2)$-manifold. Consider a pair (B, H), consisting of a set of disjoint oriented $(n+1)$-disks $B = \{B_i : D^{n+1} \to M\}$ and a collection H of oriented 1-handles $H_j : D^1 \times D^n \to M$, with $H_j(\{0,1\} \times D^n)$ embedded in $\cup_{B_i \in B} B_i(\partial D^{n+1})$, such that the orientations agree, and such that for each i, j, the components of $H_j((0,1) \times D^n) \cap B_i(D^{n+1})$ are a finite collection of n-disks embedded in $H_j((0,1) \times D^n)$, such that each component separates $H_j(\{0\} \times D^n)$ from $H_j(\{1\} \times D^n)$. This implies that $H_j((0,1) \times D^n) \cap B_i(D^{n+1})$ is contained in the interior of the image of B_i; such intersections between a handle and a base are called *ribbon intersections* or *ribbon singularities*. A handle can intersect any number of bases in ribbon intersections, and it can intersect the same base multiple times (or zero). We will in general identify the maps B_i and H_j with their images. The disks in B are called *bases*, while the elements of H are called *(fusion) handles* or *bands*. Let K be the link defined as $K = (\cup_{B_i \in B}(\partial B_i - \cup_{H_j \in H} H_j)) \cup (\cup_{H_j \in H} H_j(D^1 \times \partial D^n))$. Then K is called a *ribbon link*, and the pair (B, H) are a *ribbon presentation* for K. We identify two ribbon presentations, (B, H) and (B', H'), of K if there is a path between them in the space of ribbon presentations of K. This is equivalent to requiring that $|B| = |B'|$, $|H| = |H'|$, and that there exists an ambient isotopy f_t of M, $t \in [0,1]$, such that f_0 is the identity, and for all $B_i \in B, H_j \in H, B'_i \in B', H'_j \in H'$, $f_1 \circ B_i = B'_i$, and $f_1 \circ H_j = H'_j$.

The subspace $R = (\cup_{B_i \in B} B_i) \cup (\cup_{H_j \in H} H_j)$ is called a *ribbon solid* for the ribbon presentation or for the link. Note that a given R may be a ribbon solid for multiple ribbon presentations. If an n-link K has a ribbon presentation then it is called a *ribbon link*.

Observe that since we require that the handles be glued to the bases in an orientation-preserving manner, K will be orientable. Note also that for ribbon 1-links, the components of a ribbon link may not be ribbon knots. However, for



FIGURE 1. Two simply equivalent ribbon presentations with distinct ribbon intersections along a single handle. Here, the bases are indicated with shading; ribbon intersections are shown by thickened lines.

n-links with $n \geq 2$, each component of a ribbon link will automatically be a ribbon knot.

Given a ribbon link K, there are in general multiple ribbon presentations for K, making it convenient to consider notions of equivalence for ribbon presentations. We will, for the moment, restrict our attention to knots; the generalization to links is straightforward. Let K be a ribbon knot with ribbon presentations (B, H) and (B, H'). These presentations are called *simply equivalent* if $|H| = |H'|$, $|B| = |B'|$, and there exists an isotopy f_t, $t \in [0, 1]$, of M, such that f_0 is the identity, and for all $B_i \in B, H_j \in H, B'_i \in B', H'_j \in H'$, $f_1 \circ B_i|_{\partial D^{n+1}} = B'_i|_{\partial D^{n+1}}$, and $f_1 \circ H_j = H'_j$.

Note that simple equivalence may create new intersections of handles with the interiors of bases, as illustrated in Fig. 1.

Lemma 2. *For $n \geq 2$ a ribbon presentation (B, H) in a simply-connected manifold M is determined uniquely up to simple equivalence by specifying the following ribbon data:*
(a) the cardinality of B and,
(b) for each $H_j \in H$, a sequence of bases, $B_{j1}, ..., B_{jk}$, through which the handle H_j passes through, up to reversal of that sequence.

Proof. Clearly there is a ribbon presentation fulfilling every possible ribbon data as above. We need to show that such ribbon presentation is unique, up to simple equivalence. Note that for for each j, the sequence $B_{j1}, ..., B_{jk}$, determines the homotopy class of the handle H_j in $M - \cup_i \partial B_i$ uniquely. Since $n \geq 2$ and the core of the handle is 1-dimensional, the homotopy class of the handle determines its isotopy class as well. □

In $n = 1$ this is of course not the case, since the homotopy class of the handle does not determine its isotopy class in $M - \cup_i \partial B_i$. In addition, one must specify a cyclic ordering of handles glued to a base for $n = 1$.

There is a notion of equivalence between ribbon presentations called *stable equivalence*, defined by Nakanishi, [7]. This is the equivalence relation generated by simple equivalence together with the following three moves and their inverses, which are illustrated in Fig. 2:



FIGURE 2. The three moves permitted for ribbon presentations under stable equivalence.

(1) Add a new base B_0 with a handle H_0 starting on B_0 and terminating on any other base, without passing through any bases.
(2) *Handle pass*: Isotope H_j through $H_{j'}([0,1]\times(D^n - \partial D^n))$, leaving $H_j|_{\{0,1\}\times D^n}$ fixed.
(3) *Handle slide*: If H_j connects B_i with $B_{i'}$, and $H_{j'}$ has an end on B_i, we may slide that end along H_j over to $B_{i'}$.

Observe that for a knotted n-sphere, $|H| = |B| - 1$ by Euler characteristic arguments. It is easy to check that for $n \geq 2$, a ribbon n-knot K will have the homeomorphism type of an n-sphere with $|H| - |B| + 1$ 1-handles attached.

Given a ribbon n-knot K, it is an open question whether any two ribbon presentations for K are related by stable equivalence for $n \geq 2$. For the case $n = 1$, Nakanishi and Nakagawa provide a family of knots, each of which admits multiple ribbon presentations which are not stably equivalent, [8, 7].

3. Quandles

A *quandle*, [3], (also called a *distributive groupoid* in [6]) $(Q, *)$ consists of a set Q and a binary operation $*$ on Q such that for all $a, b, c \in Q$, the following equalities hold.

(1) $a * a = a$.
(2) There is a unique $x \in Q$ such that $x * b = a$.
(3) $(a * b) * c = (a * c) * (b * c)$.

When the operation $*$ is implied, we may denote the quandle by Q, suppressing the operation.

In general, it is common to denote the $*$ operation in any quandle Q by using the conjugation notation, that is, for any quandle we may define $a^b = a*b$. Analogously, we will denote the element x stipulated by property (2) by $a^{\bar{b}}$, i.e. $(a^b)^{\bar{b}} = a$. By convention, we interpret $a^{bc} = (a^b)^c$.

Quandles as knot invariants were introduced in [3, 6]. Here we recall the definition of a quandle associated to a knot, as given by Joyce for classical knots in [3]. The geometric definition holds for any codimension-2 knot, however.

Let L be a link in M, with L and M assumed to be orientable, and let $N(L)$ be an open tubular neighborhood of L. Then $M - N(L)$ is a manifold with boundary containing $\partial\overline{N(L)}$. The fundamental group of this space is called the *link group*. We will follow the convention that the product of two homotopy classes of paths goes from right to left, i.e. $\gamma'\gamma$ is the homotopy class of a path which follows γ and then γ'. We call the image P of the homomorphism induced by the injection $\partial\overline{N(L)} \to M - N(L)$ the *peripheral* subgroup of $\pi_1(M - L)$. Note that P is defined only up to conjugation; any of its conjugates are also peripheral subgroups.

A meridian m of L is the boundary of the disk fibre of the tubular neighborhood of L considered as a disk bundle over L. Note that the orientations of M and of L determine an orientation of m. A meridian is defined uniquely up to conjugation.

The definition of the quandle, and of the fundamental group of a link complement, requires concatenation of paths. We will use the following convention for path concatenation: given two paths γ, γ', we will write $\gamma'\gamma$ to indicate the path that follows γ first, and then follows γ'. This convention makes it easier to discuss the action of the link group on the quandle.

We define the quandle $Q(M, L)$ associated to the pair (M, L) and a chosen base point $b \in M - N(L)$ as follows. The elements of $Q(M, L)$ are homotopy classes of paths in $M - N(L)$ that start at b and end in $\partial\overline{N(L)}$, where homotopies are required to preserve these two conditions. The quandle operation, $[q]^{[q']}$, between two elements $[q], [q'] \in Q(M, L)$ is defined as the homotopy class of $q\overline{q'}mq'$, where the overbar indicates that the path is followed in reverse and m is the meridian of L passing through the endpoint of q, cf. 3. It is a straightforward exercise to show that this binary operation is well-defined and it satisfies the definition of a quandle operation. We call $Q(M, L)$ the *link quandle* of L (or the knot quandle if L is a knot).



FIGURE 3. The quandle operation for the fundamental quandle of a link L in any space M. The open circle indicates the basepoint of M, while the black circles represent points on the boundary of a tubular neighborhood of L.

Note that there is a right action of $\pi_1(M - N(L))$ on $Q(M, L)$: $[q]g$ is the equivalence class including the homotopy class of the path qg.

The following theorem was proved for classical knots in [3]. The arguments in [3] are easily generalized to any knots, as is done in [9].

Theorem 3. *For knots, $Q(M, K)$ is determined by $\pi_1(M - K)$ with its peripheral structure.*

4. Ribbon n-Knots with Isomorphic Quandles

Let us say that in a ribbon presentation (B, H) a handle $h \in H$ is a *trivial handle* if both ends are attached to the same base and h can be isotoped so it does not meet any bases except at its ends.

Definition 4. *Let K, K' be n-knots in S^{n+2}, and let $T = S^1 \times S^{n-1}$. Let T_n indicate the connected sum of n copies of T that are trivially embedded, that is, which bound embedded copies of $S^1 \times D^n$. We will say that K and K' are* weakly equivalent *if $K \# T_n$ is isotopic to $K' \# T_m$.*

Note that in [5], weak equivalence is called "stable equivalence," as gluing on T may be thought of as a stabilization of the knot. Because this term is also used to refer stable equivalence of ribbon presentations, we are adopting the term "weak equivalence" instead to avoid any confusion.

Definition 5. *Let $(B, H), (B', H')$ be ribbon presentations for n-links, $n \geq 2$. We will say that they are k-*weakly stably equivalent *if they are stably equivalent after attaching k trivial handles, and that they are* weakly stably equivalent *if they are k-weakly stably equivalent for some finite k.*

Given a ribbon presentation (B, H), the bases b_i in B are a collection of embedded $(n + 1)$ disks. We will say that a path in $S^{n+2} - K$ *passes through* a base $b_i \in B$ if it intersects the interior of b_i. Note that in general position, a path

can pass through a base positively or negatively, depending on whether it passes through it in the same direction as the normal vector or in the opposite direction.

Remark 6. *To specify a handle $h \in H$, it suffices to specify the starting base to which $\{0\} \times D^n$ is glued, the signed intersections of the handle in the order it passes through bases, and the terminal base to which $\{1\} \times D^n$ is glued. Any two handles meeting these conditions are smoothly isotopic and thus equivalent.*

Theorem 7. *Let K be a ribbon knot in S^{n+2}, $n \geq 2$, of arbitrary genus. Suppose that K admits a base and handle presentation (B, H) with k bases, and $\pi_1(S^{n+2} - K) \cong \mathbb{Z}$. Then (B, H) is $(k-1)$-weakly stably equivalent to an unknotted ribbon knot.*

Proof. Let $h_2, ..., h_k$ be $(k-1)$ new trivial handles, with both ends connected to base B_1. Perform handle slides so that h_i starts on B_1 and terminates on B_i, twisting them at the end so that they pass through bases equally many times positively as negatively. Let γ_i be a curve which travels to B_1 without passing through any bases, then follows the handle h_i to B_i (traveling through the same bases in the same order) and returns to the basepoint for the fundamental group without passing through any additional bases. Since $\pi_1(S^{n+2} - K) \cong \mathbb{Z}$, and since by construction γ is homologically trivial, γ bounds a disk in $S^{n+2} - (K \cup h_2 \cup ... \cup h_k)$. This disk may be taken to be smooth and immersed by various approximation theorems, [4, Theorem 10.16, Theorem 10.21] and [1, Theorem 2.5]. By general position, this disk meets itself at most in isolated points. Sliding h_i along this disk we may, by handle passes only, make h_i into a handle connecting B_1 to B_i without passing through any bases. Now all handles in H may be moved by handle passes so they only pass through base B_1, after which all other bases may be deleted after appropriate handle slides. But a ribbon presentation with a single base is always stably equivalent to the unknotted ribbon presentation. □

For a ribbon n-knot K with $n \geq 2$ and ribbon presentation (B, H), there is a natural presentation for the knot quandle $Q(S^{n+2}, K)$. This presentation is generated by the bases in B and has one relation for each handle $h_j \in H$. The relation r_j has the following form: let b_s and b_e be the starting and ending bases for h_j, and let h_j follow the word $w = b_1 b_2 ... b_n$ (here w is a word in the bases in B, where each base can appear positively or negatively, depending on whether the handle passes through it in a direction that agrees with the normal vector or opposite to it) as it moves from b_s to b_e. Then the relation r_j is given by the equation $b_e = b_s^{\overline{b_1}\,\overline{b_2} \cdots \overline{b_n}}$. Geometrically, the generator corresponding to a base $b \in B$ is given by an equivalence class of paths containing a path that goes from the basepoint to the base b without passing through any other bases.

Theorem 8. *A ribbon n-knot, $n \geq 2$, with ribbon presentation (B, H), has a knot quandle whose presentation is as described in the previous paragraph.*

Proof. This is proved by repeated application of the Van Kampen theorem. Alternately, it is not difficult to write down the double point curves in a projection of a ribbon knot where each base and handle is embedded under the projection. Then the double point curves correspond to handles passing through bases, and the relations may be worked out using the computations as shown in e.g. [2] (which are themselves based on the Van Kampen theorem). □

Remark 9. *Let $Q(S^{n+2}, K)$ be the quandle of a ribbon knot K with presentation (B, H). To specify an element of $Q(S^{n+2}, K)$, it suffices to specify a path γ by assuming that γ starts at the basepoint for the quandle, then specifying the ordered signed intersections of the path with the bases, and finally specifying on which base the path terminates.*

This follows from general position arguments.

Remark 10. *We will denote the generator of a quandle corresponding to a base b by using the same symbol; whether we are referring to the base or the quandle element will be clear from context.*

Throughout the following, we will make extensive use of the following fact, which forms part of the geometric definition of the quandle of a knot complement. Let K be a knot in S^{n+2}, and let x be the basepoint for the quandle $Q(S^{n+2}, K)$. Let $N(K)$ be the closure of a tubular neighborhood of K. Recall that elements of the quandle are equivalence classes of paths γ which start on x and terminate on $\partial N(K)$. Two paths γ, γ' are equivalent as quandle elements iff they are homotopic through paths starting on x and terminating on $\partial N(K)$. Therefore, there exists a disk $D_{\gamma,\gamma'}$ whose boundary consists of the union of $\gamma \cup \gamma'$ with some path on $\partial N(K)$. By various approximation theorems, [4, Theorem 10.16, Theorem 10.21] and [1, Theorem 2.5], we may assume this disk is smooth and immersed, and therefore by general position we may assume that $D_{\gamma,\gamma'}$ is immersed and embedded except at a finite number of points, provided that $n \geq 2$. In fact for $n \geq 3$, general position arguments show that the disk can be taken to be embedded. Thus, the homotopy may be taken to be a homotopy through embedded paths, that is, a smooth isotopy of the path which traces out the disk $D_{\gamma,\gamma'}$.

We will repeatedly make use of the existence of such isotopies. All our applications are essentially uses of the following construction. Let (B, H) be a ribbon presentation for K. Let h, h' be handles such that $(B, H \cup \{h\})$ and $(B, H \cup \{h'\})$ are ribbon presentations. Recall that a handle is an embedding of $D^1 \times D^n$ into S^{n+2} such that $\{0, 1\} \times D^n$ is embedded in elements of B. We will refer to $h|_{[0,1]\times\{0\}}$ as the *core* of h. Observe that once the core of h is specified, the handle itself is specified up to isotopy. Now suppose that h, h' have the property that $h|_{[0,1/2]\times\{0\}} = h'|_{[0,1/2]\times\{0\}}$. Let $\gamma = h|_{[1/2,1]\times\{0\}}, \gamma' = h'|_{[1/2,1]\times\{0\}}$. Now γ, γ' determine quandle elements for $Q(S^{n+2}, K)$.

Lemma 11. *Let K, (B, H), h, h', γ, γ' be as described in the previous paragraph, and suppose that $\gamma \cong \gamma'$ as elements of $Q(S^{n+2}, K)$. Then there is a stable equivalence between $(B, H \cup \{h\})$ and $(B, H \cup \{h'\})$ which involves only performing handle slides isotoping h to agree with h'.*

Proof. Let $D_{\gamma,\gamma'}$ be the homotopy disk giving the equivalence of γ, γ' as elements of $Q(S^{n+2}, K)$. This is a disk in the complement of K. This disk may be perturbed so that its interior does not meets h or h', by general position arguments. Now $D_{\gamma,\gamma'}$ intersects itself only in isolated points. Therefore, we may slide $h|_{[1/2,1]\times\{0\}}$ along this disk until it agrees with $h'|_{[1/2,1]\times\{0\}}$. During this process we will be sliding the end of the core of h around on $\partial N(K)$ so we may perform some handle slides. □

Remark 12. *This argument cannot be extended to an argument for an analogous result for handles h and h' that are already part of the ribbon presentation for K.*

This is because in that case, the disk $D_{\gamma,\gamma'}$ might pass through the interior of the handle h or h' themselves. This cannot happen in the above case because γ, γ' are homotopic using a homotopy in the complement of K, and, therefore, the homotopy cannot move the path through the interior of h or h' since they are not handles for the presentation of K.

Lemma 13. *Let K be a ribbon n-knot, $n \geq 2$, with ribbon presentation (B, H). Suppose that the quandle $Q(S^{n+2}, K)$ is generated by $B - \{b\}$. Then, by adding a trivial handle to (B, H), we can merge b and some $b' \in B - \{b\})$ into a single base through stable equivalence.*

Proof. Let h be the new trivial handle, with both ends on b. We may isotope h so that the midpoint of the core of h passes through the basepoint x for the quandle while it remains a trivial handle. Now the path that goes from the basepoint to b is equivalent in the quandle to a path γ from the basepoint to some other base b' which does not pass through the base b. There is a quandle homotopy disk connecting these two paths which does not intersect h (since all the relations in the quandle are geometrically given by the handles in H). We may slide the portion of h between its midpoint and terminal point along this disk so that it joins b and b' and follows the path γ, using the method described in Lemma 11. Since γ does not pass through b, we can now perform handle passes through h so that any handles passing through b are moved to instead pass through b'. Then the two bases can easily be joined. □

Corollary 14. *Every ribbon n-knot K, $n \geq 2$, with ribbon presentation (B, H), is $(|B| - 1)$-weakly stably equivalent to a ribbon n-knot with presentation (B', H'), where $B' \subseteq B, H \subseteq H'$, with the property that for all $b \in B'$, the subquandle generated by $B' - \{b\}$ is a proper subquandle of $Q(S^{n+2}, K)$.*

Theorem 15. *Let K, K' be ribbon n-knots, $n \geq 2$, with ribbon presentations $(B, H), (B, H')$, respectively, of the same genus, such that the identity map on B induces a quandle isomorphism between $Q(S^{n+2}, K)$ and $Q(S^{n+2}, K')$. Then they are $|H|$-weakly stably equivalent.*

Proof. Let $h' \in H'$ connect b_0, b_1. Attach a trivial handle h to H with both ends on b_0. Let h be isotoped so that the midpoint of its core passes through the basepoint x for the quandle. The path from x to b_0 following the core of h and the path from x to b_1 following h' are homotopic in the quandle $Q(S^{n+2}, K)$; thus we may slide h's latter half along the homotopy disk, as described in Lemma 11. After this is done, h will pass through exactly the same bases as h'. Repeat this for each handle in H'. Then perform the same operation using the trivial handles we have added to H', modifying them so that all the handles originally in H are now in H' as well. We now have all the handles in H in H' as well, and all the handles in H' have been added to H. Thus we have modified the two ribbon presentations so they have identical sets of bases and handles. □

Theorem 16. *Let K, K' be ribbon n-knots, $n \geq 2$, with ribbon presentations of the same genus $(B, H), (B', H')$, respectively. Let $f : B \to Q(S^{n+2}, K')$ be a function which extends to a quandle isomorphism $Q(S^{n+2}, K) \to Q(S^{n+2}, K')$. Then K, K' are $(2|B'| + |H|)$-weakly stably equivalent with bases given by B and $f(B)$ respectively.*

Proof. For each $b_i \in B$, we obtain a quandle element $b_i' = f(b_i) \in Q(S^{n+2}, K')$. The b_i' are paths from the basepoint to some base $B_i' \in B'$. Create a trivial base B_i'' connected by a handle to B_i' by a handle that does not pass through any bases. Then we may slide b_i' so that it terminates on B_i''. Clearly b_i' does not pass through B_i''. Therefore, we can pull B_i'' back along b_i', contracting the quandle element, until we have so isotoped B_i'' such that b_i' is now a path from the basepoint to B_i'' which does not pass through any bases.

Now the collection of B_i''s generates the quandle $Q(S^{n+2}, K')$. By Lemma 13, (B', H') is $|B'|$-weakly stably equivalent to a knot whose bases consist of exactly the B_i''s. Add an equal number of handles to H. Then by Theorem 15 the result follows. □

Theorem 17. *Let K, K' be ribbon n-knots (of the same genus) with ribbon presentations $(B, H), (B', H')$, respectively. Suppose the knot quandles $Q(S^{n+2}, K)$ and $Q(S^{n+2}, K')$ are isomorphic. Then K and K' are $(2|B| + |H|)$-weakly stably equivalent.*

Proof. Let $f : Q(S^{n+2}, K) \to Q(S^{n+2}, K')$ be the isomorphism. Then f defines a map from B to $Q(S^{n+2}, K')$ defined by sending $b \mapsto f(b)$ (where we send the base to the quandle element which its standard generator is sent to by f. Now apply Theorem 16. □

Corollary 18. *Let K, K' be as in the previous theorem, but with possibly different genus. Then they are weakly stably equivalent.*

Corollary 19. *Let $(B, H), (B', H')$ be two ribbon presentations for a single n-knot K, $n \geq 2$. Then they are $(2|B| + ||H|)$-weakly stably equivalent.*

This is related to the question posed in [7], where it is asked whether two ribbon presentations for the same n-knot with $n \geq 2$ are necessarily stably equivalent. In [8] an infinite family of examples are constructed showing that two ribbon presentations for the same 1-knot do not need to be stably equivalent. In fact examples are constructed of 1-knots with n stably inequivalent ribbon presentations for any $n \in \mathbb{N}$.

References

[1] M. Adachi, *Umekomi to hamekomi (Embeddings and immersions)*, translated by K. Hudson, AMS (1993).

[2] J.S. Carter, S. Kamada, and M. Saito, *Surfaces in 4-space*, Springer (2004).

[3] D. Joyce, *A Classifying Invariant of Knots, the Knot Quandle*, Journal of Pure and Applied Algebra **23** 37-65 (1982).

[4] J. Lee, *Introduction to smooth manifolds*, Springer (2006).

[5] C. Livingston, *Stably irreducible surfaces in S^4*, Pacific Journal of Mathematics, Vol. 116, No. 1 (1985).

[6] S. Matveev, S.V. *Distributivnye grupoidy v teorii uzlov*, Mat. Sbornik **119** 1 (1982) 78-88 (in Russian). English Version: *Distributive groupoids in Knot Theory*, Mat. Sbornik **47** 73-83 (1984).

[7] Y. Nakanishi, *On ribbon knots, II*, Kobe. J. Math 3 (1986) 77-85.

[8] Y. Nakanishi, Y. Nakagawa, *On ribbon knots*, Math. Sem. Notes Kobe Univ. 5 (1982) 423-430.

[9] B. Winter, *Virtual, wlded, and ribbon links in arbitrary dimensions*, Ph.D. dissertation, SUNY Buffalo (2014).